

UNITED STATES DEPARTMENT OF COMMERCE National Oceanic and Atmospheric Administration

NATIONAL MARINE FISHERIES SERVICE
Habitat Conservation Division
James J. Howard Marine
Sciences Laboratory
74 Magruder Road
Highlands, New Jersey 07732

March 10, 2000

Mr. Kevin P. Madden, Director
Office of Pipeline Regulation
Federal Energy Regulatory Commission
888 First Street, N.E., Room 1A
Washington, D.C. 20426

Re: OPR/DEER/ERC II, Millennium Pipeline Company, L.P. Docket No. CP98-150-000, § 375.3

Dear Mr. Madden:

This letter is in response to your Environmental Information Request (EIR) regarding potential construction windows in Haverstraw Bay for the subject project. Since the EIR was issued, we have discussed the sensitivity of aquatic resources that use Haverstraw Bay with the New York State Departments of State (DOS), Coastal Management Program (NYCMP) and Environmental Conservation (DEC). We focused on seasonal windows that are appropriate for construction activities proposed in Haverstraw Bay, and are in general agreement with New York State on appropriate dredging windows for Haverstraw Bay. Our conclusion is that a fall working window is preferred. As requested in the EIR, we have appended a list of references used to evaluate suspended sediment impacts and their relationship to resources that use Haverstraw Bay. Copies of specific documents or contact information for individuals consulted are available upon request. Finally, the ecological sensitivity of Haverstraw Bay remains the basis of our opposition to the proposed Millennium Pipeline Company alignment.

Haverstraw Bay has been recognized by NYCMP as one of the most important fish and wildlife habitats in the Hudson River Estuary. This is formally recognized by New York's designation of the area as a Significant Coastal Fish and Wildlife Habitat pursuant to New York State Law and the Federal Coastal Zone Management Act. NOAA concurs with the designation. Many aquatic resources of concern, notably anadromous, estuarine-dependent and certain marine species use and are dependent upon this habitat for critical functions and values, notably spawning, nursery, feeding and overwintering activities. These resources are managed under a variety of federal legislative actions, including the Fish and Wildlife Coordination Act, the Magnuson-Stevens Fishery Conservation and Management Act, the Emergency Striped Bass Act, Coastal Zone Management Act and the Endangered Species Act.

The NY Coastal Fish and Wildlife rating form and narrative for the state designation of Haverstraw Bay habitat details the basis for this designation, and includes the NYCMP's conclusion that this habitat is irreplaceable. We have and continue to concur with the NYCMP analysis and findings. Since the New York State Department of State already has provided FERC the documentation for site designation, we have not appended a copy to this communication.

Maintenance dredging of all but the smallest projects for Haverstraw Bay should be conducted only between September 1 and November 15 of any calendar year. However, new dredging does not meet the habitat impairment test criteria established by the NYCMP for this site, and we cannot support any construction window for new work. And, as our staff already has indicated to FERC and the project proponents, the selected alignment and installation technique would produce unacceptable and avoidable impacts to aquatic resources including endangered and other special concern species. Instead, an out-of-Bay, less damaging alignment should be pursued if a crossing of the Hudson River is necessary or appropriate. We will evaluate new proposals for crossings outside of the Haverstraw Bay habitat that the project proponents find technically feasible. Those alternatives may include rerouting the pipeline to a more northerly alignment after it exits Bowline power station, accessing Bowline power station with a lateral connection, or terminating the pipeline at Bowline power station and eliminating the Hudson River crossing altogether.

In Haverstraw Bay, aquatic resources would be especially impacted from mid-November through August. Overwintering fish would be vulnerable between approximately November 15 and March 31, and an April through August dredging restriction is needed to protect sensitive age groups of fishery resources. We do not expect that an EFH assessment for this area would conclude that additional seasonal restrictions beyond those indicated above are necessary. However, other mitigative measures may be required for work scheduled in the September-mid November window. Any such measures under consideration should be discussed in the FERCs analysis of direct, indirect and cumulative impacts, referencing the appropriate species/lifestage(s) to which benefits would accrue. Similarly, if it is FERC's preliminary determination that no additional measures would be necessary, the EFH assessment should state how that conclusion was reached.

With respect to shortnose sturgeon and how we considered our suggested maintenance dredging period for Haverstraw Bay, the topic remains a subject for continued discussions among the FERC, USACE and NMFS. The intent would be to minimize interactions with that species during concentration periods. By limiting maintenance dredging to September through Mid-November, some key habitat uses by shortnose sturgeon would be protected. However, some level of interaction would occur in the proposed maintenance window and must be addressed as part of the Section 7 consultation. We will review the need to adjust the proposed window or make other recommendations or stipulations upon receipt of the Biological Assessment. As indicated in previous correspondence, it is premature for us to make a final

determination regarding shortnose sturgeon, especially since some level of take, as defined by the ESA, appears unavoidable.

I hope that this information is helpful. If you or your staff wishes to discuss this information further, please contact Diane Rusanowsky or Michael Ludwig at (203) 579-7004.

Sincerely,

Stanley W. Gorski

Field Offices Supervisor

cc: COE/NY - Heidi Firstencel (Albany Field Office)

USF&WS - Diane Mann-Klager (Cortland Field Office)

USEPA - Dan Montella (Region 2) NYSDOS - Steven Resler (NYCMP) NYSDEC - Margaret Duke (Region 3)

NMFS, PRD - Mantzaris

eirnewry - 001

Selected References

- Able, K.W. and M.P. Fahay. 1998. The First Year in the Life of Estuarine Fishes in the Middle Atlantic Bight. Rutgers University Press, New Brunswick, NJ.pp342.
- Auld, A.H. and J.R. Schubel. 1978. Effects of Suspended Sediment on Fish Eggs and Larvae: A Laboratory Assessment. Estuar. And Coastal Mar Sci. vol:6, 153-164.
- Bisbal, G.A. and D.A. Bengtson. 1995. Effects of Delayed Feeding on Survival and Growth of Summer Flounder (Paralicthys dentatus) larvae. Mar. Ecol. Prog. vol:121:1-3, pp301-306.
- Blaxter, J.H.S. 1974. The Early Life History of Fish The Proceedings of an International Symposium Held at the Dunstaffnage Marine Research Lab. of the Scottish Marine Biological Assoc. at Oban, Scotland, 17 -23 May, 1973. Springer-Verlag, NY, pp765.
- Bohlen, W.F., M.M. Howard-Strobel, D.R. Cohen and E.T. Morto. 1996. An Investigation of the Dispersion of Sediments Resuspended by Dredging Operations in New Haven Harbor. Contribution 112, DAMOS, US Army Corps of Engineers, New England Division, Concord, MA.
- Bokuniewicz, H.J., J. Gebert, R.B. Gordon, J.L. Higgins, P. Kaminsky, C.C. Pilbeam, M. Reed, C. Tuttle. 1978. Field Study of the Mechanics of the Placement of Dredged Material at Open-Water Disposal Sites. Technical Report D-78-7. US Army Waterways Experiment Station. Vicksburg, MS.
- Breitburg, D.L. 1988. Effects of Turbidity on Prey Consumption by Striped Bass Larvae. Trans Amer. Fish. Soc. vol:117, pp 72-77.
- Buckel, J.A., D.O. Conover, N.D. Steinberg and K.A. McKown. 1999. Impact of Age-0 Bluefish (Pomatomus saltatrix) Predation on Age-0 Fishes in the Hudson Estuary:

 Evidence for Density-dependent Loss of Juvenile Striped Bass (Morone saxatilis). Can. J. Fish. Aquat. Sci. vol:56, pp275-287.
- Burton, W.H. 1993. Effects of Bucket Dredging on Water Quality in the Delaware River and the Potential for Effects on Fisheries Resources. Prepared for Delaware Basin Fish and Wildlife Management Cooperative. Versar, Inc. Columbia, MD.
- Chesney, E.J.,Jr. 1989. Estimating the Food Requirements of Striped Bass Larvae Morone saxatilis: Effects of Light, Turbidity and Turbulence. Mar. Ecol. Prog. Ser. vol:53 pp191-200.

- Collins, M.A. 1995. Dredging-Induced Near-Field Resuspended Sediment Concentrations and Source Strengths. Miscell. Paper D-95-2. US Army Corps of Engineers, W.E.S., Vicksburg, MS.
- Gibson, M. Personal Communication.
- Hartman, K.J. 1997. Progress Report: Seasonal and Spatial Assessment of Bay Anchovy Population Abundance: Biology and Biometrics in the Hudson River. Submitted to Hudson River Foundation, NY, NY.
- Johnston, D.D. and D.J. Wildish. 1982. Effect of Suspended Sediment on Feeding by Larval Herring (Clupea harengus harengus L.). Bull Environm. Contam. Toxicol. vol:29, 261-267.
- Laurence, G.C. 1976. A Bioenergetic Model for the Analysis of Feeding and Survival Potential of Winter Flounder (Pseudopleuronectus americanus) During the Period from Hatching to Metamorphosis. ICES Council Meeting 1976 collected papers.
- Leonard, J.B.K. and S.D. McCormick. 1999. The Effects of Migration Distance and Timing on Metabolic Enzyme Activity in an Anadromous Clupeid, the American shad (Alosa sapidissima). Fish Physio. & Biochem.. vol:20:2 pp 163-179
- Limburg, K.E., M.L. Pace, D. Fischer, and K.K. Arend. 1997. Consumption, Selectivity, and Utilization of Zooplankton by Larval Morone Spp. In a Seasonal Pulsed Estuary. Trans.AFS vol:126 pp607-621.
- Limburg, K.E., M.L. Pace, and K.K. Arend. 1999. *Growth, Mortality and Recruitment of Larval Morone Spp. In Relation to Food Availability and Temperature in the Hudson River.* Fish. Bull. vol:97 pp80-91.
- McGovern, J.C. and J.E. Olney. 1996. Factors Affecting Survival of Early Life Stages and Subsequent Recruitment of Striped Bass on the Pamunkey River, Virginia. Can. J. Fish. Aquat. Sci. vol: 53 pp1713-1726.
- McLellan, T.N., R.N. Havis, D.F. Hayes and G.L. Raymond. 1989. Field Characteristics of Sediment Resuspension Characteristics of Selected Dredges. Tech. Report HL-89-9, US Army Corps of Engineers, WES, Vicksburg, MS.
- May, R.C. 1974. Larval Mortality in Marine Fishes and the Critical Period Concept. Contribution No. 437, Hawaii Institute of Marine Biology, University of Hawaii, Kaneohe, HI. Pp16.
- Messiah, S.N., D.J. Wildish and R.H. Peterson. 1981. *Possible Impact of Sediment Dredging and Spoil Disposal on the Miraichi Bay Herring Fishery*. Can. Tech. Rept. Fish. Aquatic Sci. No. 1008.

- Minello, T.J.,R.J. Zimmerman and R.J. Martinez. 1987. Fish Predation on Juvenile Brown Shrimp,
 Penaeus aztecus Ives: Effects of Turbidity and Substratum on Predation
 Rates. US Fish. Bull. vol:85, pp59-70.
- Morgan, R.P., V.J. Rasin and L.A. Noe. 1973. Effects of Suspended Sediments on the Development of Eggs and Larvae of Striped Bass and White Perch. Natural resources Institute, Chesapeake Biological Laboratory, University of Maryland, Center for Environmental and Estuarine Studies. Pp20.
- Mourente, G., A. Rodriques, A. Grau, & E. Pastor. 1999. *Utilization of Lipids by Dentex dentex L. Larvae During Lecitotropha and Subsequent Starvation.* Fish Physio. & Biochem. vol:21:1, pp45-58.
- Nelson, D. Personal communications. NOAA/NMFS/NEFSC, Milford, CT.
- Nelson, D.A. and J.L. Wheeler. 1997. The Influence of Dredging-Induced Turbidity and
 Associated Contaminants Upon Hatching Success and Larval Survival of
 Winter Flounder, Pleuronectus americanus, A Laboratory Study. Final Report
 on Grant CWF #321-R. Submitted to State of CT Depart. Of Environ.
 Protection. Hartford, CT. Pp. 57
- Newcombe, C.P. and D.D. MacDonald. 1991. Effects of Suspended Sediments on Aquatic Ecosystems. North American J. of Fish. Man. vol:11 pp72-82.
- Pereira, J. Personal communications. NOAA/NMFS/NEFSC, Milford, CT.
- Profiles Research and Consulting Groups, Inc. 1980. Seasonal Restrictions on Dredging Projects by NMFS in the Northeast. Prepared for USDOC/NOAA/NMFS, Gloucester, MA.
- Rusanowsky, D. D.A. Nelson and M. Ludwig. *Impacts of Dredging: A Winter Flounder Case Study.* Section 6, Habitat Impacts: Special Topics. USDOC, NOAA, NMFS, Gloucester, MA.pp42.
- Schubel, J.R. and J.C.S. Wang.1973. The Effects of Suspended Sediment on the Hatching Success of Perca flavescens (Yellow Perch) Morone americana (White Perch), Morone saxatilis (Striped Bass), and Alosa pseudoharengus) (Alewife) Eggs. Chesapeake Bay Institute, Johns Hopkins University Special Report No 30, Ref. 73-3, pp77.
- Schultz, E.T. and D.O. Conover. 1997. Latitudinal Differences in Somatic Energy Storage:

 Adaptive Responses to Seasonality in an Estuarine Fish (Atherinidae:

 Menidia menidia). Oecologia. vol: 109, pp 516-529.

- Schultz, E.T., D.O. Conover and A. Ehtisham. 1998. The Dead of Winter: Size-Dependent Variation and Genetic Differences in Seasonal Mortality Among Atlantic Silverside (Atherinidae: Menidia menidia) from Different Latitudes. Can. J. Fish. Aquat. Sci. vol: 55:5 pp 1149-1157.
- Sullivan, B.K. and D. Hancock. 1977. Zooplankton and Dredging: Research Perspectives from a Critical Review. Water Resources Bull. vol:13:3 pp461-468.
- Tavolaro, J. R. 1984. A Sediment Budget Study of Clamshell Dredging and Ocean Disposal Activities in the New York Bight. Environmental Geology and Water Sciences. vol. 6(3):133-140.